

AD-A182 947

SUM RULES FOR OPTICAL EXTINCTION AND SCATTERING BY
SMALL PARTICLES(U) PENNSYLVANIA STATE UNIV UNIVERSITY
PARK DEPT OF METEOROLOGY C F BOHREN 18 AUG 86
AFOSR-TR-87-0873 AFOSR-84-0145


1/1

UNCLASSIFIED

F/G 4/1

ML





DTIC FILE COPY

(2)

REPORT DOCUMENTATION PAGE

AD-A182 947

HIC
EGTE
20 1987

1b. RESTRICTIVE MARKINGS

3. DISTRIBUTION/AVAILABILITY OF REPORT

Approved for public release,
distribution unlimited
unlimited

2a. DECLASSIFICATION/DOWNGRADING SCHEDULE

4. PERFORMING ORGANIZATION REPORT NUMBER

5. MONITORING ORGANIZATION REPORT NUMBER(S)

AFOSR-TR-87-0873

6a. NAME OF PERFORMING ORGANIZATION

Pennsylvania State University

6b. OFFICE SYMBOL

(If applicable)

7a. NAME OF MONITORING ORGANIZATION

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH

6c. ADDRESS (City, State and ZIP Code)

Meteorology Department
University Park Pennsylvania 16802

7b. ADDRESS (City, State and ZIP Code)

Bldg 410
Bolling AFB DC 20332-6448

8a. NAME OF FUNDING/SPONSORING ORGANIZATION

AFOSR

8b. OFFICE SYMBOL

(If applicable)

NE

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

AFOSR-84-0145

8c. ADDRESS (City, State and ZIP Code)

Bldg 410
Bolling AFB DC 20332-6448

10. SOURCE OF FUNDING NOS.

PROGRAM
ELEMENT NO.61102F
2306PROJECT
NO.2306
C1TASK
NO.

CH

WORK UNIT
NO.

11. TITLE (Include Security Classification)

Sum Rules for Optical Extinction and Scattering

by Small Particles

12. PERSONAL AUTHOR(S)

Craig F. Bohren

13a. TYPE OF REPORT

Final

13b. TIME COVERED

FROM 16 May 84 TO 15 May 86

14. DATE OF REPORT (Yr., Mo., Day)

10 August 86

15. PAGE COUNT

9

16. SUPPLEMENTARY NOTATION

17. COSATI CODES

FIELD GROUP SUB. GR.

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

We have established that the sum rule for extinction is not changed when magnetic dipole as well as electric dipole radiation is considered. Moreover, the limitation of electron mean free paths in metallic particles does not invalidate sum rules. When metallic particles are sufficiently small, the effect of their boundaries is to decrease peak extinction while simultaneously broadening it in such a way that integrated extinction remains constant.

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT

UNCLASSIFIED/UNLIMITED ☒ SAME AS RPT ☐ DTIC USERS ☐

21. ABSTRACT SECURITY CLASSIFICATION

UUUU

UNCLASSIFIED

22a. NAME OF RESPONSIBLE INDIVIDUAL

Major Joseph Hager

22b. TELEPHONE NUMBER
(Include Area Code)

202-7674933

22c. OFFICE SYMBOL

Electroptical
NE Material Science

**Sum Rules for Optical Extinction and Scattering
by Small Particles**

Final Report

Contract No. AFOSR-84-0145

Craig F. Bohren

**Department of Meteorology
Pennsylvania State University
University Park, Pennsylvania 16802**

10 August 1986

Submitted to

**Major Joseph Hagar
Air Force Office of Scientific Research
Directorate of Electronic and Material Science
Bolling Air Force Base, Washington DC 20332-5260**

Abstract of First Year Effort

A central problem in developing effective smoke obscurants for space applications is to maximize volumetric extinction or absorption at the wavelength of the incident electromagnetic radiation. Although this is a problem without a general solution, it is possible to make recommendations based on very general considerations. One can establish bounds not on extinction at any wavelength, but rather extinction integrated over all wavelengths.

A sum rule for extinction is a bound on integrated volumetric extinction. The existence of such sum rules requires suitable behavior of volumetric extinction at high and low frequencies. If, in the limit of low frequency, extinction decreases as the $1 + \sigma$ power of frequency, where σ is any positive number, then a sum rule exists. At low frequencies, the extinction cross section of an ellipsoid composed of either a simple free-electron metal or a simple insulator is proportional to frequency squared, which ensures the existence of a sum rule for extinction by such particles. Only a single type of sum rule exists because of the rather stringent requirements on both the low- and high-frequency behavior of particle extinction cross sections.

We have established that the sum rule for extinction is not changed when magnetic dipole as well as electric dipole radiation is considered. Moreover, the limitation of electron mean free paths in metallic particles does not invalidate sum

rules. When metallic particles are sufficiently small, the effect of their boundaries is to decrease peak extinction while simultaneously broadening it in such a way that integrated extinction remains constant.

Although integrated extinction is independent of particle size, sum rules imply that the smallest particles will have the greatest volumetric extinction. When a particle is illuminated by an electromagnetic wave, various electromagnetic modes are excited. The larger the particle, the greater the number of modes. Thus, a fixed amount of total extinction must be shared by modes, the number of which increases with increasing particle size. This implies that the smallest particles have the greatest extinction.

Two classes of materials, free-electron metals and polar insulators, with vastly different properties, nevertheless have in common that integrated extinction by particles of these materials is approximately the same.

Volumetric extinction of 10^4 cm^{-1} should be looked upon as a kind of lower bound: it is the value that can be obtained without much-effort. What is the upper bound? We must look to small (compared with the wavelength) metallic particles for the greatest possible volumetric extinction. Fewer modes are excited in a sphere than in a needle or a disk. Peak extinction by a small aluminum sphere is about 10^7 cm^{-1} . We claim that this is a practical upper bound. Moreover, this bound cannot

<input checked="" type="checkbox"/>
<input type="checkbox"/>
<input type="checkbox"/>
des
or



A-1

be obtained over a wide range of frequencies. A practical upper bound over at least one decade from the ultraviolet to the near infrared is 10^6 cm^{-1} , and this would have to be obtained with a mixture of particles having different shapes.

Abstract of Second Year Effort

One of the essential lessons to be learned from sum rules is that extinction cannot be arbitrarily specified. Integrated volumetric extinction is independent of particle size and depends on only the low frequency behavior of the material of which the particle is composed. This in turn implies that extinction at one frequency is related to extinction at other frequencies. From this idea another related one evolved: the scattering coefficients are not independent. A proof of this was constructed for scattering by a sphere. The Mie coefficients are completely specified by the first four of them. A proof of this and its physical interpretation is given in Appendix A, a manuscript on recurrence relations for Mie scattering coefficients, which will appear in a future issue of Journal of the Optical Society of America.

Appendix A

Recurrence Relations for the Mie Scattering Coefficients

to appear in

Journal of the Optical Society of America

February 1987

Recurrence Relations for the Mie Scattering Coefficients

Craig F. Bohren

Department of Meteorology
Pennsylvania State University
University Park, Pennsylvania 16802

Abstract

The Mie scattering coefficients satisfy recurrence relations: a_{n-1} , b_{n-1} , a_n , and b_n , determine a_{n+1} and b_{n+1} . It is therefore possible, in principle, to generate the entire set from the first four, which has a simple interpretation. Each term in a multipole expansion of an electrostatic field can be obtained by differentiating the preceding term. The Mie coefficients are terms in a multipole expansion of a particular electromagnetic field, that scattered by an arbitrary sphere. By analogy, it is not surprising that all these coefficients can be generated from the electric and magnetic dipole and quadrupole terms. Moreover, the recurrence relations for the Mie coefficients contain finite differences, in analogy with the infinitesimal differences (derivatives) in the multipole expansion of an electrostatic field.

The solution to the problem of scattering by an arbitrary sphere (Mie theory) yields an infinite set of coefficients¹⁻⁵, which depend on various Bessel functions and their derivatives. The usual approach to computing these coefficients is to compute the Bessel functions by recurrence⁵⁻⁸. This might be called the natural or direct way of computation. It seems so natural, so obvious, that, to my knowledge, no one has ever done otherwise, although there are hints in Ref. 4 (p. 51). What is usually at issue is not if one should compute Bessel functions but rather how to do it in the fastest and most accurate way. I have posed this question differently. The Bessel functions in the expressions for the scattering coefficients satisfy recurrence relations. Furthermore, Verner⁹ showed that the numerators and denominators of these coefficients satisfy recurrence relations. Thus, is it not likely that the coefficients themselves satisfy recurrence relations? In the following paragraphs I show that they do and give a physical interpretation of why. My notation and conventions are those of Ref. 5.

The Mie scattering coefficients can be written as follows:

$$a_n = \frac{x D_n(mx) \psi_n(x) + mn \psi_n(x) - mx \psi_{n-1}(x)}{x D_n(mx) \xi_n(x) + mn \xi_n(x) - mx \xi_{n-1}(x)}, \quad (1)$$

$$b_n = \frac{mx D_n(mx) \psi_n(x) + n \psi_n(x) - x \psi_{n-1}(x)}{mx D_n(mx) \xi_n(x) + n \xi_n(x) - x \xi_{n-1}(x)}, \quad (2)$$

where D_n is the logarithmic derivative ψ'_n/ψ_n , and ψ_n, ξ_n are Riccati-Bessel functions. The size parameter x is $2\pi a/\lambda$, where a is the radius of the sphere and λ is the wavelength of the

incident light; m is the refractive index of the sphere relative to that of the surrounding medium. The logarithmic derivative satisfies the recurrence relation

$$D_{n-1} = \frac{n}{mx} - \frac{1}{D_n + n/mx}, \quad (3)$$

and both of the Riccati-Bessel functions satisfy the recurrence relation

$$z_{n-1} = \frac{2n+1}{x} z_n - z_{n+1}. \quad (4)$$

From Eqs. (1)-(4) it follows that any set of six coefficients are given by

$$a_{n-1} = \frac{uy + A_3uz + B_1y + A_4z}{uv + A_3uw + B_1v + A_4w}, \quad (5)$$

$$b_{n-1} = \frac{uy + B_1y + A_2z}{uv + B_1v + A_2w}, \quad (6)$$

$$a_n = \frac{uy + A_1y + A_2z}{uv + A_1v + A_2w}, \quad (7)$$

$$b_n = \frac{uy + B_1y + B_2z}{uv + B_1v + B_2w}, \quad (8)$$

$$a_{n+1} = \frac{uy + A_5uz + A_6y + A_7z}{uv + A_5uw + A_6v + A_7w}, \quad (9)$$

$$b_{n+1} = \frac{uy + B_3y + A_2z}{uv + B_3v + A_2w}, \quad (10)$$

where $u = D_n$, $v = \xi_n$, $w = \xi_{n-1}$, $y = \psi_n$, $z = \psi_{n-1}$ and

$$A_1 = \frac{mn}{x}, \quad A_2 = -m, \quad B_1 = \frac{n}{mx}, \quad B_2 = -1/m,$$

$$A_3 = \frac{n(1 - m^2)}{m^2 x}, \quad A_4 = \frac{n^2 - m^2 x^2 - m^2 n^2}{m^3 x^2},$$

$$A_5 = \frac{x(m^2 - 1)(n + 1)}{(n + 1)(2n + 1)(1 - m^2) + m^2 x^2},$$

$$A_6 = \frac{nm^2 x + (n + 1)^2 (2n + 1)(m^2 - 1)}{mx\{(n + 1)(2n + 1)(1 - m^2) + m^2 x^2\}},$$

$$A_7 = \frac{(n + 1)^2 (1 - m^2) - m^2 x^2}{m\{(n + 1)(2n + 1)(1 - m^2) + m^2 x^2\}},$$

$$B_3 = \frac{(2n + 1)m^2 - (n + 1)}{mx}.$$

Four quantities (e.g., u , v/y , w/y , z/y) determine any of the five coefficients a_{n-1} , b_{n-1} , a_n , b_n , a_{n+1} (or b_{n+1}). Thus, it is possible to solve for any one of these coefficients (e.g., a_{n+1} or b_{n+1}) in terms of the other four. To do so, we may consider u to be fixed in Eqs. (5)-(8) and write them in the form

$$\begin{pmatrix} a_n(A_1 + u) & a_n A_2 & -(A_1 + u) & -A_2 \\ b_n(B_1 + u) & b_n B_2 & -(B_1 + u) & -B_2 \\ b_{n-1}(B_1 + u) & b_{n-1} A_2 & -(B_1 + u) & -A_2 \\ a_{n-1}(B_1 + u) & a_{n-1}(A_3 u + A_4) & -(B_1 + u) & -(A_3 u + A_4) \end{pmatrix} \begin{pmatrix} v \\ w \\ y \\ z \end{pmatrix} = 0 \quad (11)$$

The determinant of the coefficient matrix in Eq. (11) must vanish if its solution is to be nontrivial. By setting this determinant equal to zero, one obtains a cubic equation satisfied by u . One of its factors is $B_1 + u$, which follows from inspection of Eq. (11); the root $u = -B_1$ is obviously not the one desired. The correct value of u is one of the roots of the quadratic equation

$$u^2 - \left(A_1 \Delta + \frac{A_2 - A_4}{A_3} \right) u + \frac{A_1 (B_2 - A_4)}{A_3} \Delta = 0, \quad (12)$$

where

$$\Delta = \frac{(a_n - b_n)(a_{n-1} - b_{n-1})}{(b_n - a_{n-1})(a_n - b_{n-1})}. \quad (13)$$

To obtain these equations it was also necessary to use the relations $A_2 B_2 = 1$ and $A_1 B_2 = A_2 B_1$. Which of the two roots of Eq. (12) is the correct one must be determined by some auxiliary condition (e.g., the magnitude of the scattering coefficients cannot exceed unity). Given this root, one can substitute it in Eq. (11) and determine any three of the set (v, w, y, z) as multiplicative functions of the fourth. When these functions and u are substituted into Eq. (9), a_{n+1} is obtained as a function of a_{n-1} , b_{n-1} , a_n , b_n , the index n , and the properties of the sphere (x and m). Similarly, b_{n+1} is so obtained. This completes the proof that any set of four consecutive scattering coefficients determine the following two.

This result has a simple interpretation. It has long been

known that each term in the expansion of an electrostatic field can be obtained by differentiation of the preceding term. This idea was developed by Maxwell^{10,11}, who used the term "points" of the n th order, whereas in modern work they would be referred to as " 2^n poles" or "multipoles" (see, e.g., Ref. 12). The Mie coefficients are terms in a multipole expansion of a particular electromagnetic field, that scattered by an arbitrary sphere. By analogy, it is not surprising that all of these coefficients can be generated from the electric (a_1) and magnetic (b_1) dipole terms and the electric (a_2) and magnetic (b_2) quadrupole terms. Moreover, the recurrence relations for the Mie coefficients contain finite differences (see Eq. 13), in analogy with the infinitesimal differences (derivatives) in the multipole expansion of an electrostatic field. A possible interpretation of why the multipoles in Mie theory are connected by finite differences is that this theory applies to particles with finite (relative to the wavelength) dimensions. In contrast, the wavelength is infinite (alternatively, the speed of light is infinite) in electrostatics. A further difference between electrostatics (and magnetostatics) and electrodynamics is that in static theories the electric and magnetic terms are distinct: electric terms are connected only to electric terms (and similarly for magnetic terms). But when the propagation time over characteristic distances is not negligible compared with the period (e.g., when $x = \omega a/c$ is not negligible, where c is the speed of light, and a is the radius of a particle illuminated by light of circular frequency ω), then electric and

magnetic multipoles are inextricably connected.

Whether or not the recurrence relations can simplify scattering calculations must be determined by extensive calculations. It is not known if the recurrence relations are stable, either upward or downward. And one must devise a simple test for which root of Eq. (12) is the proper one. These are matters for further investigation. Nevertheless, merely knowing that the scattering coefficients can be generated by recurrence gives new insights. It is for this reason that I have set them down, not because they will necessarily reduce computation time.

I am indebted to Nels Shirer, Robert Wells, Warren Wiscombe, and Akhlesh Lakhtakia for their valuable suggestions and comments. And I sincerely thank William Doyle and Milton Kerker not only for helping me with the technical aspects of the work reported here but also for encouraging me to carry it out. This work was supported, in part, by a grant from the Air Force Office of Scientific Research, Directorate of Electronic and Material Science.

References

1. J. A. Stratton, Electromagnetic Theory (McGraw-Hill, New York, 1941), Sec. 9.25.
2. H. C. van de Hulst, Light Scattering by Small Particles (Wiley, New York, 1957), Ch. 9.
3. M. Born and E. Wolf, Principles of Optics, 3rd. ed. (Pergamon, Oxford, 1965), Sec. 13.5.
4. M. Kerker, The Scattering of Light and Other Electromagnetic Radiation (Academic, New York, 1969), Chs. 3 and 4.
5. C. F. Bohren and D. R. Huffman, Absorption and Scattering of Light by Small Particles (Wiley-Interscience, New York, 1983), Ch. 4.
6. J. V. Dave, Subroutines for Computing the Parameters of the Electromagnetic Radiation Scattered by a Sphere, IBM Order Number 360D-17.4.002 (1968).
7. W. J. Lentz, "Generating Bessel functions in Mie scattering calculations using continued fractions," Appl. Opt. 15, 668 (1976).

8. W. J. Wiscombe, "Improved Mie scattering algorithms," Appl. Opt. 19, 1505 (1980).
9. B. Verner, "Note on the recurrence between Mie's coefficients," J. Opt. Soc. Am. 66, 1424 (1976).
10. J. C. Maxwell, A Treatise on Electricity and Magnetism, Vol. 1 (Dover, New York, 1954), p. 196.
11. E. W. Hobson, The Theory of Spherical and Ellipsoidal Harmonics (Cambridge University Press), pp. 129-132.
12. W. K. H. Panofsky and M. Phillips, Classical Electricity and Magnetism (Addison-Wesley, Reading, Mass., 1962), p. 15.

END

9-87

Dtic